

## Specific Example of How In Vitro Assays Have Been Used at RJRT as Part of a Toxicological Evaluation Program

### Eclipse

R J Reynolds has developed a cigarette that primarily heats tobacco. Under the FTC puffing regimen, the mainstream smoke from this cigarette is reduced significantly in a number of chemical compounds when compared with other cigarettes of similar "tar" delivery (Borgerding *et al.*, 1998). The mainstream particulate matter (CSC) has been evaluated in the neutral red cytotoxicity, Ames, SCE and chromosome aberration assays, and the whole smoke (vapor phase plus particulate) has been evaluated in the SCE and neutral red cytotoxicity assay (Bombick, B. *et al.*, 1998; Bombick D. *et al.*, 1998). The genotoxic potential of cigarette smoke condensate from cigarettes that primarily heat tobacco is significantly reduced compared to that of cigarettes that burn tobacco, evidenced by significant reduction in the induction of Ames bacterial mutagenicity, chromosomal aberrations, and sister chromatid exchanges. The data from these studies are presented in Figures 3 through 10.

### Summary

- RJRT has actively pursued the development of cigarettes with the potential to reduce risks by:
  1. Continuing to pursue general and specific smoke constituent reductions in tobacco-burning cigarettes.
  2. Continuing to develop and refine alternative cigarette designs.
  3. Finding ways to alter "tar"/nicotine ratio.
  4. Ensuring no product modifications add to the biological activity.
- RJRT has developed and applied assays to evaluate the progress of cigarette modifications.
- A regulatory framework should be designed that allows free-market forces to develop products that have the potential to reduce risks.
- A tiered testing strategy for toxicological evaluation has been used at RJRT to determine the level and degree of testing required which include chemical and biological assays.
- Short-term in vitro assays are a subset of the biological assays which have been developed or applied by RJRT with success for use on cigarette smoke and cigarette smoke condensate.

Boutwell, R.K. Model systems for defining irritation, promotion, and progression of skin neoplasms. In Slaga, T.J., Klein-Szanto, A.J.P., Boutwell, R.K., Steven Som, D.E., Spitzer, H.L., and D'Motta, B., Skin Carcinogenesis: Mechanisms and Human Relevance. Alan R. Liss, Inc. New York, 3, 1987.

Boutwell, R.K., Urbach, F., and Carpenter, G. Chemical carcinogenesis: B, Experimental models. In Lacrum, D.D. and Iverson, D.H. (editors) Biology of Skin Cancer (Excluding Melonomas). International Union Against Cancer, Geneva, 109, 1981.

Bowden, D.H. Macrophages, dust and pulmonary diseases. Exp. Lung Res. 12, 89, 1987.

Bremier, L.H. Molecular mechanisms of oxygen radical carcinogenesis and mutagenesis: The role of DNA base damage. Molecular Carcinogen. 3, 188, 1990.

Brody, A.R. Asbestos exposure as a model of inflammation inducing interstitial pulmonary fibrosis. In Gallin, Goldstein, and Snyderman (editors) Inflammation, 2<sup>nd</sup> edition. Raven Press, New York, 1033, 1981.

Brusick, David. in Principles and Methods of Toxicology, ed. By A. Wallace Hayes. Raven Press (New York), 545, 1994.

Bus, J.S. and Gibson, J.E. Paraquat: Model for oxidant inhaled toxicity Environ. Health Perspect. 55, 37, 1984.

Butterworth, B.E. and Goldsworthy, T.L. The role of cell proliferation in multistage carcinogenesis. Cell Proliferation. 198 (2), 683, 1991.

Cervitti P. Prooxidant states and tumor promotion, Science. 227, 375, 1985.

Cheeseman, K.H. Albano, E.F., Tomasi, A., and Slater, T.F. Biochemical studies on the metabolic activation of halogenated alkane. Environ Health Perspect. 64, 85, 1985.

Chichester, C.H. Philpot, R.M. Weir, A.J., Buckpitt, A.R. and Plopper, C.G. Characterization of the cytochrome P-450 monooxygenase system in nonciliated bronchial epithelial (Clara) cells isolated from mouse lung. Am. J. Respir. Cell Molec. Biol. 4, 179, 1991.

Clothier, R.H., Hulme, L.M., Smith, M., and Balls, M. Comparison of the in vitro cytotoxicities and acute in vivo toxicities of 59 chemicals. Mol. Toxicol. 1, 571, 1989.

Coggins, C.R.E., Doolittle, D.J., Lee, C.K., Ayres, P.H., Mosberg, A.T., Bolin, D.C., Burger, G.T., and Hayes, A.W. Histopathology, urine mutagenicity and bone marrow cytogenetics of mice exposed nose-only to smoke from cigarettes which burn or heat tobacco. Inhalation Toxicology 2, 407, 1990.

Bombick, D.W., Bombick, B.R., Ayres P.H., Putnam, K., Avalos, J., Borgending, M.F., and Doolittle D.J. Evaluation of the genotoxic and cytotoxic potential of mainstream whole smoke and smoke condensate from a cigarette containing a novel carbon filter. Fund. Appl. Toxicol. 39, 11, 1997.

Bombick, D.W. and Doolittle, D.J. The effect of cell type on chemical toxicity in the neutral red cytotoxicity assay. The Toxicologist. 12(1), 373, Abstract 1460, 1992.

Bombick, D.W. and Doolittle, D.J. The role of chemical structure and cell type in the cytotoxicity of low-molecular-weight aldehydes and pyridines. In Vitro Toxicology. 8(4), 349, 1995.

Bombick, D.W., Putnam, K., Bombick, B.R., Morgan, W.T., Ayres, P.H., and Doolittle D.J. In vitro assessment of biological activity of mainstream smoke from cigarettes which burn or primarily heat tobacco. The International Toxicologist. 7, 73, 1995.

Bombick, D.W., Putnam, K., Bombick, B.R., Morgan, W.T., Ayres, P.H. and Doolittle, D.J. In vitro assessment of biological activity of mainstream smoke from cigarettes which burn or primarily heat tobacco. The Toxicologist. 30(1), 129, 1996.

Bombick, D.W., Putnam, K., and Doolittle, D.J. Comparative cytotoxicity studies of smoke condensates from different types of cigarettes and tobaccos. Toxicology In Vitro 12, 241, 1998.

Bombick, D.W. Putnam, K., and Doolittle, D.J. The contribution of several hydroxy- and dihydroxybenzenes found in cigarette smoke to the over all cytotoxicity of the cigarette smoke mixture. Poster presented at the INVITOX international meeting, September 14-18, 1998.

Bond, J.A., Johnson, N.F., Snipes, M.B., and Mauderly, J.L. DNA adduct formation in rat alveolar type II cells: Cells potentially at risk for inhaled diesel exhaust. Environ. Molec. Mutagen. 16, 64, 1990.

Border, W.A. and Noble, N.A. Transforming growth factor beta in tissue fibrosis. New England J. of Medicine. 331 (9), 1286, 1994.

Borgending, M.F., Bodner, J.A., Chung, H.L., Morgan, P.P., Morrison, C.C., Risner, C.H., Rogers, J.C. Summons, D.F., Uhrig, M.S., Wendelboe, F.N., Wingate, D.E., and Winkler, L.S. Chemical and biological studies of a new cigarette that primarily heats tobacco. Part I. Chemical composition of mainstream smoke. Food Chem. Toxicol. 36, 169, 1998.

Domin, B.A., Devereux, T.R., and Philpot, R.M. The cytochrome P-450 monooxygenase system of rabbit lung: enzyme components, activities, and induction in the nonciliated bronchial epithelial cell alveolar type II cell, and alveolar macrophage. Molec. Pharmacol. 30, 296, 1986.

Doolittle, DJ, CA Rahn, GT Burger, CK Lee, B. Reed, E. Riccio, G Howard, GT Passananti, ES Vesell, and AW Hayes. Effect of cooking methods on the mutagenicity of food and on urinary mutagenicity of human consumers. Food Chem. Tox. 27 (10), 657, 1989.

Doolittle, DJ, CK Lee, JL Ivett, JC Mirsalis, E Riccio, C Rudd, GT Burger, and A W Hayes. Genetic toxicology studies comparing the activity of sidestream smoke from cigarettes which burn or only heat tobacco. Mutat. Res. 249:59, 1990a.

Doolittle, DJ, CK Lee, JL Ivett, JC Mirsalis, E Riccio, CJ Rudd, GT Burger, AW Hayes. Comparative studies on the genotoxic activity of mainstream smoke condensate from cigarettes which burn or only heat tobacco. Environ Mol. Mutagen. 15:93, 1990b.

Driscoll, K.E. Carter, J. M., Howard, B.W., and Hassenbein, D.G. Mutagenesis in rat lung epithelial cells after in vivo silica exposure or ex vivo exposure to inflammatory cells. Am. J. Respir. Crit. Care Med. 149, A553, 1994.

Driscoll, K.E. Maurer, J.K. Lindenschmidt, R.C. Romberger, D. Rennard, S.I., and Crosby, L. Respiratory tract responses to dust: Relationships between dust burden, lung injury, alveolar macrophage, fibronectin release, and the development of pulmonary fibrosis. Toxicol. Appl. Pharmacol. 106, 88, 1990.

Dzau, V.J., Pathobiology of Atherosclerosis and plaque complications. American Heart Journal. 128 (6 Pt.2) 1300, 1994.

Ekwall, B., Bondesson, I., Castell, J.V., Gomez-Lechon, M.J., Hellberg, S. Hogberg, J., Jover, R., Ponsada, X., Romert, L., Stenberg, K., and Walum, E. Cytotoxicity Evaluation of the first ten MEIC chemicals: acute lethal toxicity in man predicted by cytotoxicity in five cellular assays and by oral LD50 tests in rodents. ATLA, 17,83, 1989.

Ekwall, B., Clemendson, D., Barile, F., Calleja, M.C., Castell, J., Chesme, C. Clothier, R., Curreu, R., Dierictx, P., McFarlane-Abdullah, E., Ferro, M., Fiskejo, G., Garza-Ocanas, L., Gomez-Lechan, M.J., Gulden, M., Harvell, J.D., Isamaa, B., Vamus, J., Kerszman, G., Kristen, V., Kunimoto, M., Karenl'ampi, S., Lavrijsen, K., Lewan, L., Ohno, T., Persone, G., Roguet, R., Ramert, L., Sawyet, T., Seibert, H., Shrivastava, R., Stadtlamder, K., Stammati, A., Tamaka, K., Triglia, D., Valentino, M., Walum, E., Wang, X., Zveco, F., and Maiback, H. Comparison between human skin irritancy and in vitro cytotoxicity from 77 systems for the first 12 MEIC chemicals. In Vitro Tox. 7, 156, 1994.

Faotrel, A., Chesne', C., Guillozo, A., DeSousa, G., Placidi, M., Rhamani, R., Braut, F., Picliou, J., Hoellinger, H., Vintezous, P., Diarte, I., Melcion, C., Cordier, A., Larenzon, G., Benincourt,

- Cohen, C. Dossou, K.G., Rougier, A. and Roguet, R. Measurement of inflammatory mediators produced by human keratinocytes in vitro: a predictive assessment of cutaneous irritation. Toxicol. In Vitro. 5, 407, 1991.
- Claxton, L.D., Allen, J, Auletta, A, Mortelmans, K, Nestmann, E., and Zeiger, E. Guide for the *Salmonella typhimurium*/mammalian microsome tests for bacterial mutagenicity. Mutat. Res. 189, 83, 1987.
- Colton, C.A. and Zakhari, S. Role of free Radicals in alcohol-induced tissue injury. In Basken, S.I. and Salem, H. (editors) Oxidants Antioxidants and Free Radicals, Taylor and Francis, Washington, DC. 259, 1997.
- Cross, C.E. Oxygen radicals and human diseases. Am. Intern. Med. 107, 526, 1987.
- Crystal, R.G., Bitterman, P.B., Remard, S.L, Hance, A., and Keogh, B.A. Interstitial lung disease of unknown cause: Disorders characterize by chronic inflammation of the lower respiratory tract. N. Engl. J. Med. 310, 154, 1984.
- Curvall, M., T. Jansson, B. Pettersson, A. Hedin, and C. Enzell. (1985). In vitro studies of biological effects of cigarette smoke condensate. I. Genotoxic and cytotoxic effects of neutral, semivolatile constituents. Mutat. Res. 157, 169, 1985.
- Curvall M, and C. Enzell. (1984). Evaluation of the biological activity of cigarette smoke condensate fractions using six *in vitro* short-term tests. J. Toxic. Environ. Health 14:163, 1984.
- De Flora, S., R. Balansky, L. Gasparini, and A. Camoirano Bacterial mutagenicity of cigarette smoke and its interaction with ethanol. Mutagenesis 10 (1), 47, 1995.
- DeMarini, D.M. Genotoxicity of tobacco smoke and tobacco smoke condensate. Mutat. Res. 114, 59, 1983.
- Devereux, T.R. Serabjit-Singh, C., Slaughter, S.R., Wolf, C.R. Philpot, R.M., and Fouts, J.R. Identification of cytochrome P-450 isozymes in nonciliated bronchial epithelial (Clara) and alveolar type II cells isolated from rabbit lung. Exp. Lung Res. 2, 221, 1981.
- Devereux, T.R. Alveolar type II and Clara cells; isolation and xenobiotic metabolism. Environ. Health Persp. 56, 95, 1984.
- Digiovanni, J., Kruozewski, and Chemicek, K.J. Studies on the skin tumor promoting actions of Chrysarobin in Butterworth, B.E. and Slaga, T.J., Banbury Report 25: Nongenotoxic Mechanisms And Carcinogenesis Cold Spring Harbor Laboratory, p 25-39, 1987.
- Dixit, R., Khanna, A.K., Waseem, M., Dagra, S, and Kaw, J.L. Alterations in in vitro functional activities of alveolar macrophages exposed in vivo to mineral dusts. Vet. Hum. Toxicol. 32, 517, 1990.

- M., Vannier, B., Favreux, R., Peloux, A.F., Bichet, N., Govy, D., and Cano, J.D. A multicentre study of acute in vitro cytotoxicity in rat liver cells. Toxicology in Vitro, 5 (5/6), 543, 1991.
- Fedullo, A.J. and Ettensohn, D.B. Bronchoalveolar lavage in lymphangitic spread of adenocarcinoma to the lung. Chest, 87, 129, 1985.
- Fels, A.O.S. and Cohn, Z.A. The alveolar macrophage. J. Appl. Physiol. 60, 353, 1980.
- Floyd, R.A., The role of 8- hydroxyguanine in carcinogenesis. Carcinogenesis, 11(9), 1447, 1990.
- Forkert, P.G. Vessey, M.L., Park, S.S., Gelborm, H.V., and Cole, S.P.C. Cytochrome p-450 in murine lung. An immunohistochemical study with monoclonal antibodies. Drug Metab. Disp. 17, 551, 1989.
- FRAME. 1991. Animals and alternatives in toxicology: Present status and future prospects (The Second Report of the FRAME Toxicity Committee). Toxic Subst. 8, 403.
- Frazier, J.M. Scientific perspectives on the role of in vitro toxicity testing in chemical safety evaluation. In Jolles, G. and Cordier, A. (editors) In Vitro Methods in Toxicology. Academic Press, London. 99, 522, 1992.
- Gailit, J., and Clark, R.A.F. Wound repair in the context of extracellular matrix. Current Opinion in Cell Biology, 6 (5), 717, 1994.
- Gordner, D.E., and Graham, J.A. Increased pulmonary disease mediated through altered bacterial defenses. In Sanders, C.L., Schneider, R.P. Doyle, G.E., and Ragar, H.A. (editors) Pulmonary Macrophage and Epithelial Cells. Proceedings Sixteen Annual Hanford Symposium Series. Richland, Washington, 1, 1976.
- Graham, J.A. and Gardner, D.E. Immunotoxicity of air pollutants. In Dean, J.H., Luster, M.I., and Munson, A.E. (editors) Immunotoxicology and Immunopharmacology. Raven Press, New York, 367, 1985.
- Gueniche, A. and Ponc, M. Use of human skin cell cultures for the estimation of potential skin irritants. Toxicol. In Vitro, 7, 15, 1993.
- Guyton, K.Z., and Kensler, T.W. Oxidative mechanisms in carcinogenesis. Br. Med. Bull. 49 (3), 523, 1993.
- Halliwell, B. Reactive oxygen species in living systems: Source, biochemistry and role in human disease. Am. J. Med. 92 (supplement 3C), 145, 1992.
- Harris, J.O. Swenson, E.W., and Johnson, J.E. Human alveolar macrophages: comparison of phagocytic ability, glucose utilization, and ultrastructure in smokers and non smokers. J. Clin. Invest. 69, 2086, 1970.

Haslet, A. and Henson, P.M. Resolution of inflammation . In Clark R.A.F. and Henson P.M. (editors) The Molecular and Cellular Biology of Wound Repair. Plenum, New York, 185, 1988.

Hoffmann, G.R. Genetic toxicology. In Casarett and Doull's Toxicology, Fifth edition, ed. By Klaassen, C. New York: McGraw-Hill, 269, 1996.

Hunninghale, G.W., Gadek, J.E., Kawanami, O., Ferraus, V.J. and Crystal, R.G. Inflammatory and immune process in the human lung in health and disease; evaluation by bronchoalveolar lavage. Am. J. Pathol. 97, 149, 1979.

Hutton, J. and C. Hackney. Metabolism of cigarette smoke condensates by human and rat homogenates to form mutagens detectable by *Salmonella typhimurium* TA1538. Cancer Research 35, 2461, 1975.

IARC (International Agency for Research on Cancer). Monographs on the evaluation of the carcinogenic risk of chemicals to humans: Tobacco smoking 38, 168, 1986.

Jansson, T., M. Curvall, A. Hedin, and C. Enzell. In vitro studies of biological effects of cigarette smoke condensate. II. Induction of sister chromatid exchanges in human lymphocytes by weakly acidic, semivolatile constituents. Mutat. Res. 169, 129, 1986.

Jones, K.G., Holland, J.F., and Fouts, J.R., Benzo(a)pyrene hydroxylase activity in enriched population of Clara cells and alveolar type II cells from control and B-naphthoflavone pretreated rats. Cancer Res. 42, 4658, 1982.

Josephy, P.D., P. Gruz, and T. Nohmi. (1997). Recent Advances in the construction of bacterial genotoxicity assays. Mutat. Res. 386, 1, 1997.

Kamp, D.W., Graceffa, P., Weitzman, S.A., and Pryor, W.A. The role of free radicals in asbestos-induced diseases. Free Radical Biol. Med. 12, 293, 1992.

Kensler, T.W. Egner, P.A. Taffe, B.G. and Trush, M.A. Role of free radicals in tumor promotion and progression. In Slaga T.J., Klein Szanto A.J.P., Boutwell, R.K., Stevenson, D.E., Spitzer, H.L., and D'Motto, B. (editors) Skin Carcinogenesis: Mechanisms and Human Relevance. Alan R. Liss, Inc., New York, 223, 1989.

Kier, L. E. Yamasaki, and B.N. Ames. Detection of mutagenic activity in cigarette smoke condensate. Proc. Nat. Acad. Sci. USA 71 (10), 4159, 1974.

Klaasen, C.D. and Plaa, G.L. Comparison of the biochemical alterations elicited in livers from rats treated with carbon tetrachloride, chloroform, 1, 1, 2- trichloroethane and 1,1,1-trichloroethane. Biochem Pharmacol. 18, 2019, 1969.

Knox, P., Uphill, P.F., Fry, J.R., Benford, J., and Balls, M. The FRAME multicentre project on in vitro cytotoxicology. Food Chem. Toxicol. 24, 457, 1986.

Mizusaki, S., T. Takashima and K. Tomaru. Factors affecting mutagenic activity of cigarette smoke condensate in *Salmonella typhimurium* TA1538. Mutat. Res. 48, 29, 1977.

Nathan C.F., Murray, H.W. Cahn, Z.A. Current Concepts: The macrophages as an effector cell. New Engl. J. Med. 303, 622, 1980.

Nachman, R.L. and Silverstein R. Hypercoagulable states. Annals of Internal Medicine. 119 (8), 819, 1993.

Oasting, R.S., Van Bree, I. Van Iwaarden, J.F., Van Galde, L.M.G., and Verhoff, J. Impairment of phagocytic functions of alveolar macrophages by hydrogen peroxide. Am. J. Physiol. 259, 87, 1990.

O'Brien, K.D. and Chait, A. The biology of the artery wall in atherogenesis. Medical Clinics of North America 78 (1), 41, 1994.

OConnell, J.F., Ratsterin, J.B., and Slaga, T.J. Tumor promotion and progression in the mouse skin model In Butterworth, B.E. and Slaga, T.J. (editors) Banbury Report 25, Nongenotoxic mechanisms in carcinogenesis. Cold Spring Harbor Laboratory, 11, 1987.

Ohno, T., Gordon, D., Sam, H., et al. Gene therapy for vascular smooth muscle cell proliferation after arterial injury. Science. 265, 781, 1994.

Organization for Economic Cooperation and Development. Scientific criteria for validation of in vitro toxicity tests. OECD Environment Monograph No. 36. OECD: Paris, France, 1990.

Osborne, R., Perkins, M.M., and Roberts, D.A., Development and ultralaboratory evaluation of an in vitro human cell-based test to aid ocular irritancy assessments. Fundam. Appl. Toxicol. 28, 139, 1995.

Pham, S.H. and Thrall, R.S. (editors) Pulmonary Fibrosis, Marcel Dekker, New York, 1995.

Plaa, G.L. Free-radical-mediated liver injury. In Wallace, K.B. (editor) Free Radical Toxicology, Taylor and Frances, Washington, DC, 175, 1997.

Putnam, K.P., Bombick, D.W., Avalos, J.T., and Doolittle, D.J. Evaluation of commercially available smoke flavorings used in foods. Environ. Mol. Mutagen. 29 (Supplement 28), 42, 1997.

Putnam, K.P., Bombick, D.W., and Doolittle, D.J. Evaluation of the eight in vitro assays for assessing the cytotoxicity of cigarette smoke condensate. Environ. Mol. Mutagen. 31 (Supplement 29), 37, 1998.

Reed, B.A., C.K. Lee, B.G. Brown, C.R.E. Coggins, A.W. Hayes, and D.J. Doolittle. Cytogenetic studies in laboratory animals exposed by inhalation to mainstream smoke or



Kovacs, E.J. Fibrogenic cytokines: The role of immune mediators in the development in fibrosis. Immunol. Today. 12, 17, 1991.

Kroll, J., Bagley, A.C., Mullenbach, G.T., Hallewell, R.A., Lynch R.E. Superoxide mediates the toxicity of paraquat for cultured mammalian cells. J. Biol. Chem. 263, 1910, 1988.

Kurose, I., Higuchi, H., Kato, s., Miura, S., and Ishii, H. Ethanol- induced oxidative stress in the liver. Alcohol Clin. Exp. Res. 20 (Supplement 5), A77, 1996.

Lafi A, and J.M. Parry. The effects of ventilation and the dilution of smoke upon the clastogenic and aneugenic activity of tobacco particulate matter in cultured mammalian cells. Mutat. Res. 264, 51, 1991.

Lee, C.K., Brown, BG, Reed, E.A.; Lowe, G.D., McKarns, S.C., Fulp, C. W., Coggins, C.R.E., Ayres, P.H., Doolittle D.J. Analysis of cytogenetic effects in bone marrow cells of rats subchronically exposed to smoke from cigarettes which burn or only heat tobacco. Mutat. Res. 240: 251, 1990a.

Lee, CK, DJ Doolittle, BT Burger, and AW Hayes. Comparative genotoxicity testing of mainstream whole smoke from cigarettes which burn or heat tobacco. Mutat. Res. 242:37, 1990b.

Li, A.P. An in vitro lung epithelial cell system for evaluating the toxic potential of inhaled material. Food Chem. Toxicol. 24, 527, 1986.

Li, A.P., Hahn, F.F., Zamora, P.O., Shimizv, R.W., Henderson, R.F., Brooks, A.L., and Richards, R. Characterization of a lung epithelial cell strain with potential applications in toxicological studies. Toxicol. 27, 257, 1983.

Luger, T.A., Kock A., Danner, M., Colot, M., and Micksche, M. Production of distinct cytokines by epidermal keratinocytes. Br. J. Dermatol. 113, 145, 1985.

Maron, D. and B.N. Ames. Revised methods for the Salmonella mutagenicity test. Mutat. Res. 113, 173, 1983.

Mehendale, H.M. Commentary: Role of hepatocellular regeneration and hepatobular healing in the final outcome of liver injury. A two stage model of toxicity. Biochem. Pharmacol. 42, 1155, 1991.

Mehendale, H.M., Roth, R.A., Gaudolfi, A.E., Klaunig, J.E., Lemasteis, J.J., and Curtis, L.R. Novel mechanisms in chemically induced hepatotoxicity. FASEB Journal. 8, 128,5, 1994.

Mehra, M.R. Lavie, C.J., and Ventura, H.O. Prevention of atherosclerosis. Postgraduate Medicine. 98(1), 175, 1995.

Schaefer, E.J., Lichtenstern, A.H., Lamon-Fava, S., et al. Lipoproteins, nutrition, aging, and athero-sclerosis. American J. Clin. Nutr. 61(3 supplements), 5726, 1995.

Schulter- Hermann, R. Timmermann- Trosiener I., Barthel, G., and Bursch, W. DNA synthesis apoptosis and phenotypic expression as determinants of growth of a altered foci in rat liver during phenobarbital promotion. Cancer Res. 50, 5127, 1990.

Shaw, R.J. Benedict, S.H., Clark, R.A.F., and King, T.E. Jr. Pathogenesis of pulmonary interstitial lung disease aveolar macrophage PDGF (B) gene activation and up regulation by interferon- $\gamma$ . Am. Rev. Resp.Dis. 143, 167, 1991.

Sibilie, Y. and Reynolds, H.Y. Macrophages and polymorphonuclear neutrophils in lung defense and injury. Am.Rev. Respir. Dis. 141, 471, 1990.

Slaga, T., Klein-Szanto, A., Triplet, L. Jotti, L. Trosko, J. Skin tumor-promoting activity of benzylperoxide a widely used free-radical generator. Science. 213, 1023, 1981.

Slaga, T.J., Semar mouse skin tumorigenesis model versus other strains and stocks of mice. Environ. Health Perspect. 68, 27, 1986.

Slaga, T., Solanki, V., and Logani, M. Studies on the mechanism of action of antitumor promoting agents: Suggestive evidence for the involvement of free radicals in promotion. In Nygaard, O. and Simic, M. (editors) Radioprotectors and anticarcinogens. Academic Press, New York, 471, 1983.

Smith, C.J, SC McKarns, RA Davis, SD Livingston, BR Bombick, JT Avalos, WT Morgan, and DJ Doolittle. Human urine mutagenicity study comparing cigarettes which burn or primarily heat tobacco. Mutat. Res. 361, 1, 1996.

Sone, S., Brennan, L.M. and Creasia, W.A. In vivo and in vitro NO<sub>2</sub> exposure enhances phagocytic and tumoricidal activities of rat alveolar macrophages. J. Toxicol. Environ. Health. 11, 151, 1983.

Spiclmann, H. Ocular Irritation. In Castell, J.V. and Gomez-Lechan, M.J. (editors) In Vitro methods in Pharmaceutical Research. Academic Press, San Diego, 265, 1997.

Stadnyk, A.W. Cytokine production by epithelial cells. FASEB Journal 8 (13), 1041, 1994.

Steele, R.H, V.M. Payne, C.W. Fulp, D. C. Rees, C.K. Lee, and D.J. Doolittle. A comparison of the mutagenicity of mainstream cigarette smoke condensates from a representative sample of the US cigarette market with a Kentucky reference cigarette (K1R4F). Mutat. Res. 342, 179, 1995.

Swauger, J.E., M.J. Morton, B.B. Collie, J.T. Avalos, C.A. Rahn, E.A. Bombick, DJ Doolittle, M.F. Borgerding, and D.C. Rees. A comparison of the mainstream cigarette smoke chemistry and mutagenicity of a representative sample of the U.S. cigarette with Kentucky reference cigarettes K1R4F and K1R5F. The Toxicologist. 36 (1):152, 1997.

environmental tobacco smoke, in Chromosomal Alterations, Origin and Significance, ed. By G. Obe and A. Natarajan, Springer-Verlag (Berlin Heidelberg), 362, 1994.

Rietjens, I.M.C.M., Dormans, J.A.M.A. Rambout, R.J.A. and Van Bree, L. Qualitative and quantitative changes in cytochrome p-450 dependent xenobiotic metabolism in pulmonary microsomes and isolated Clara cell populations derived from ozone-derived rats. Toxicol. Environ. Health. 24, 51, 1988.

Roberts, W.C. Preventing and arresting coronary atherosclerosis. American Heart J. 130, 580, 1995.

Roquet, R. Cutaneous Pharmacotoxicology In vitro. In Castell, J.V., and Gomez-Lechon, M.J. (editors) In vitro methods in Pharmaceutical Research. Academic Press, San Diego, 241, 1997.

Roquet, R., Regnier, M., Cohen, C., Dossou K.G., and Rougier, A. The use of in vitro reconstructed human skin in dermatotoxicity testing. Toxicol. In Vitro. 8, 6365, 1994.

Rossi, G.P., Pavae, E., and Pessina, A.C. Role of genetic, humoral, and endothelial factors in hypertension-induced atherosclerosis J. Cardiovascular Pharmacol. 23 (Supplement 5), 575, 1994.

Rowan, A.N. and Stratman, C.J. (editors) The Use of Alternatives in Drug Research. Macmillan Press, London., 129, 1980.

Rubanji, G.M. The role of endothelium in cardiovascular homeostasis and disease. J. Cardiovascular Pharmacol. 22 (4 Supplement), 1, 1993.



Rynard, S.M. Urine Mutagenicity Assays, in Biological Markers in Epidemiology, ed. by B.S. Hulka, T.C. Wilcosky, and J.D. Griffith. New York: Oxford University Press, 56, 1990.

Saladino, A.J., Willey, J.C., Lechner, J.F., Grafstrom, R.C., LaVeck, M., and Harris C.C. Effects of formaldehyde, acetaldehyde, benzyl peroxide, and hydrogen peroxide on cultured normal human bronchial epithelial cells. Cancer Research. 45, 2522, 1985.

Samani, S.M. Husain, K., and Schlorff, E.C. Response of antioxidant system to physical and chemical stress In Baskin S.I. and Salem, H. (editors) Oxidants, Antioxidants, and Free Radicals. Taylor and Francis, Washington, DC, 125, 1997.

Sato, S., Y. Seino, T. Ohka, T. Yahagi, M. Nagao, T. Matsushima, and T. Sugimura. Mutagenicity of smoke condensates from cigarettes, cigars, and pipe tobaccos. Cancer Letters 3, 1, 1977.

Sawyer, T.W. In In vitro Cytotoxicology, Mechanisms and Methods, ed. by Frank A. Barile. CRC Press (Boca Raton), 127, 1994.



Swenberg, J.A., Gross, E.A., Randall, H.W., and Barrow, C.S. The effect of formaldehyde exposure on cytotoxicity and cell proliferation. In Clary, J.J., Gibson, J.E., and Wautz, R.S. (editors) Formaldehyde: toxicology epidemiology, mechanisms. Marcel Dekker New York, 225, 1983.

Toyokuni, S., Okamoto, K., Yodai, J., and Hiai, H. Persistent oxidative stress in cancer. FEBS Letters. 358, 1, 1995.

Trivedia, P., and Mowat, A.P. Carbon tetrachloride- induced hepatic fibrosis and cirrhosis in the developing rat: An experimental model of cirrhosis in childhood. Br. J. Exp. Pathol. 64, 25, 1983.

Tucker, J.D., Auletta A., Cimino, M., Dearfield, K., Jacobson-Kram, D., Tice, R., and A. Carrano. Sister chromatid-exchange: second report of the Gene-Tox program. Mutat. Res. 297, 101, 1993.

Tucker, J.D., and R.J. Preston. Chromosome aberrations, micronuclei, aneuploidy, sister chromatid exchanges, and cancer risk assessment. Mutat. Res. 365, 147, 1996.

Varani, J. and Ward, P.A. Activation of the inflammatory response by asbestos and silicate mineral dusts. In Wallace, K.B., Free Radical Toxicology, Taylor and Francis, Washington, D.C., 295, 1997.

Varani, J. and Ward, P.A. Mechanisms of endothelial cell injury in acute inflammation. Shock, 2, 311, 1994.

Yeadon, M. Free Radicals in Lung inflammation and environmental exposure to pollutants. In Blake, D. and Winyard, P.G. (editors) Immunopharmacology of Free Radical Species, Academic Press, London, 215, 1995.

Walf, R.K. Bond, J.A. Henderson, R.F., Harkena, J.R., and Mauderly, J.L. Pulmonary inflammation and DNA adducts in rats inhaling diesel exhaust or carbon black. Inhal. Toxicol. 2, 241, 1990.

Witschi, H. Proliferation of type II alveolar cells: a review of common responses in toxic lung injury. Toxicology. 5, 267, 1976.

Zeiger, E. Carcinogenicity of mutagens: predictive capability of the *Salmonella* mutagenesis assay for rodent carcinogenicity. Cancer Res 47, 1287, 1987.

Figure 5. Chromosome aberration assay of CSC  
without activation




Figure 6. Chromosome aberration assay of CSC with  
activation




Figure 3. Ames assay of CSC  
TA98 with metabolic activation





Figure 4. Ames Assay of CSC  
TA98 without metabolic activation









Figure 7. SCE assay of CSC without activation





Figure 8. SCE Assay of CSC with activation



**Figure 5.** Chromosome aberration assay of three cigarette smoke condensates without metabolic activation. The Chromosome aberration assay was conducted in Chinese Hamster Ovary (CHO) cells according to procedures described in Bombick, B.R. *et al.*, 1998. Cigarette Smoke Condensates (CSC) were prepared under a standard FTC puffing regimen of 35 ml puffs of two-second duration every minute. The three CSCs tested were Eclipse, a cigarette that primarily heats tobacco; 1R4F, a Kentucky Reference low 'tar' cigarette, and 1R5F, a Kentucky Reference ultra-low 'tar' cigarette. CHO cells were exposed to CSC for two hours and harvested 20 hours after initiation of treatment. Concentration is expressed as ug Total Particulate Matter (TPM) per ml of media.

**Figure 6.** Chromosome aberration assay of three cigarette smoke condensates in the presence of metabolic activation. The Chromosome Aberration assay was conducted in Chinese Hamster Ovary (CHO) cells in the presence of S9 metabolic activation according to procedures described in Bombick, B.R. *et al.*, 1998. Cigarette Smoke Condensates (CSC) were prepared under a standard FTC puffing regimen of 35 ml puffs of two-second duration every minute. The three CSCs tested were Eclipse, a cigarette that primarily heats tobacco; 1R4F, a Kentucky Reference low 'tar' cigarette, and 1R5F, a Kentucky Reference ultra-low 'tar' cigarette. CHO cells were exposed to CSC for two hours in the presence of a S9 metabolic activation system, and harvested 20 hours after the initiation of treatment. Concentration is expressed as ug Total Particulate Matter (TPM) per ml media.

**Figure 7.** SCE Assay of three cigarette smoke condensates without metabolic activation. The Sister Chromatid Exchange (SCE) assay was conducted in Chinese Hamster Ovary (CHO) cells according to procedures described in Bombick, B.R. *et al.*, 1998. Cigarette Smoke Condensates (CSC) were prepared under a standard FTC puffing regimen of 35 ml puffs of two-second duration every minute. The three CSCs tested were Eclipse, a cigarette that primarily heats tobacco; 1R4F, a Kentucky Reference low 'tar' cigarette, and 1R5F, a Kentucky Reference ultra-low 'tar' cigarette. CHO cells were exposed to CSC for approximately twenty-four hours and harvested approximately 26-29 hours after initiation of treatment. Concentration is expressed as ug Total Particulate Matter (TPM) per ml media.

## Legend

- Figure 1.** Assessment of cytotoxic potential of three cigarette smoke condensates using the Neutral Red assay. The cytotoxic potential of the smoke condensate of Eclipse (a cigarette which primarily heats tobacco) was compared with smoke condensates from low 'tar' 1R4F and 1R5F ultra-low 'tar' Reference cigarettes. Smoke condensates were prepared under a standard FTC puffing regimen of 35-ml puffs of two-second duration every minute. Concentration is expressed as ug total particulate matter per mL of cell culture medium. The cytotoxic response is expressed as a percentage of control (cells exposed to solvent vehicle). CHO (Chinese Hamster Ovary) cells were used in the cytotoxicity assessment with an exposure time of 24 hours. For complete details see Bombick, B.R. *et al.*, 1998.
- Figure 2.** Assessment of cytotoxic potential of the whole smoke from three cigarettes using the Neutral Red assay. The cytotoxic potential of whole smoke from Eclipse (a cigarette which primarily heats tobacco) was compared with the whole smoke from low 'tar' 1R4F and ultra-low 'tar' 1R5F reference cigarettes. Smoke was generated from a 30-port AMESA smoke generator under a FTC puffing regimen of 35-ml puffs of 2-second duration every minute. Concentration is expressed as the number of cigarettes per cubic meter. The cytotoxic response is expressed as a percentage of control (cells exposed to humidified room air). Chinese Hamster Ovary (CHO) cells were exposed to smoke for one hour and cytotoxicity assessed after 24 hours. For complete details see Bombick, D.W. *et al.*, 1998.
- Figure 3.** Ames assay of three cigarette smoke condensates with TA98 without metabolic activation. The Ames assay was conducted in the preincubation assay according to procedures described in Bombick, B.R. *et al.*, 1998. Cigarette Smoke Condensates (CSC) were prepared under a standard FTC puffing regimen of 35 ml puffs of two-second duration every minute. The three CSCs tested were Eclipse, a cigarette that primarily heats tobacco; 1R4F, a Kentucky Reference low 'tar' cigarette, and 1R5F, a Kentucky Reference ultra-low 'tar' cigarette. Concentration is expressed as ug Total Particulate Matter (TPM) per plate.
- Figure 4.** Ames assay of three cigarette smoke condensates with TA98 in the presence of metabolic activation. The Ames assay was conducted in the preincubation assay in the presence of S9 metabolic activation according to procedures described in Bombick, B.R. *et al.*, 1998. Cigarette Smoke Condensates (CSC) were prepared under a standard FTC puffing regimen of 35 ml puffs of two-second duration every minute. The three CSCs tested were Eclipse, a cigarette that primarily heats tobacco; 1R4F, a Kentucky Reference low "tar" cigarette, and 1R5F, a Kentucky Reference ultra-low 'tar' cigarette. Concentration is expressed as ug Total Particulate Matter (TPM) per plate.

**Figure 8.** SCE assay of three cigarette smoke condensates in the presence of metabolic activation. The Sister Chromatid Exchange assay was conducted in Chinese Hamster Ovary (CHO) cells in the presence of S9 metabolic activation according to procedures described in Bombick, B.R. *et al.*, 1998. Cigarette Smoke Condensates (CSC) were prepared under a standard FTC puffing regimen of 35 ml puffs of two-second duration every minute. The three CSCs tested were Eclipse, a cigarette that primarily heats tobacco; 1R4F, a Kentucky Reference low 'tar' cigarette, and 1R5F, a Kentucky Reference ultra-low 'tar' cigarette. CHO cells were exposed to CSC for two hours in the presence of a S9 metabolic activation system, and harvested approximately 29-31.5 hours after the initiation of treatment. Concentration is expressed as ug Total Particulate Matter (TPM) per ml media.

**Figure 9.** SCE Assay of Whole Smoke. The Sister Chromatid Exchange (SCE) assay was conducted in Chinese Hamster Ovary (CHO) cells according to procedures described in Bombick, D.W. *et al.*, 1998. Smoke was generated from a 30-port AMESA smoke generator under a FTC puffing regimen of 35 ml puffs of 2-second duration every minute. Concentration is expressed as the number of cigarettes per cubic meter. The three CSCs tested were Eclipse, a cigarette that primarily heats tobacco; 1R4F, a Kentucky Reference low 'tar' cigarette, and 1R5F, a Kentucky Reference ultra-low 'tar' cigarette. CHO cells were exposed to smoke for one hour and harvested 27.5 to 32 hours after the initiation of treatment.

**Figure 10.** Human Urinary Mutagenicity Study using TA98 5% S9. The Ames assay was conducted by the microsuspension modification according to procedures described in Smith, *et al.*, 1996. Twenty-two smokers and 12 nonsmokers were enrolled in a 4-week single crossover study. Each smoker consumed Eclipse cigarettes *ad libitum* for a week and their usual brand of cigarettes the other 3 weeks. A strictly controlled diet was administered to minimize ingestion of mutagenic protein pyrolysis products. 24-hour urine samples were collected weekly, concentrated using XAD-2 resin, and tested in the microsuspension Ames assay with TA98 in the presence of S9 metabolic activation. Mutagenicity is expressed as the number of revertants per 24 hour-urinary output.